

THE IMPACT OF BREASTS AND BRAS ON PHYSICAL ACTIVITY AMONGST WOMEN AND GIRLS: A SYSTEMATIC REVIEW AND META-ANALYSIS

Gabrielle Gilmer, BChE^{1,2}, Elaine Xu, BS³, Corinna Franklin, MD, AB, FAOA⁴, Nailah Adams, MD, MS, CAQSM, FAAFP⁵, Katherine Rizzone, MD, MPH^{3,6}

- ¹ Cellular and Molecular Pathology Graduate Program, Department of Pathology, University of Pittsburg School of Medicine, Pittsburg, Pennsylvania, U.S.A.
- ² Medical Scientist Training Program, University of Pittsburg School of Medicine, Pittsburg, Pennsylvania, U.S.A.
 - ³ University of Rochester School of Medicine, Rochester, New York, U.S.A.
 - ⁴ Department of Orthopaedics and Rehabilitation, Yale School of Medicine, New Haven, Connecticut, U.S.A.
- ⁵ Department of Family Medicine, University of North Carolina Chapel Hill School of Medicine, Chapel Hill, North Carolina, U.S.A.
 - ⁶ Department of Orthopaedics and Pediatrics, University of Rochester Medical Center, Rochester, New York, U.S.A.

BACKGROUND: Breast motion may cause breast pain in active women, which can negatively impact physical activity. Sports bras are essential pieces of sporting equipment and were designed to help women comfortably participate in sports and engage in physical activity. Thus, the purpose of this systematic review was to summarize how breasts and bras influence physical activity in women and girls.

METHODS: A systematic review was conducted by performing electronic searches of PubMed, Physiotherapy Evidence Database (PEDro), Cumulative Index to Nursing and Allied Health Literature (CINAHL), and Cochrane Central Register of Controlled Trials. Title, abstract, and full text screening were performed by two independent reviewers to identify articles investigating how breasts or bras impact physical activity related outcomes in women or girls. We performed meta-analyses to evaluate the effect of (i) breast size, (ii) bra type, and (iii) breast reduction on (i) breast pain, (ii) physical activity level, and (iii) breast biomechanics. Standardized mean differences and pooled standard deviations of outcome measures were calculated using the DerSimonian-Laird method. All meta-analyses were performed using SPSS Statistics Software.

RESULTS: Of the 323 articles identified, 77 were included in the final analysis. The majority of articles studied participants who were 20-29 years old, identified as white, lived in Western countries, were nonathletes, and weighed between 45-65 kg (BMI: 18.5-29.9), while they were performing a running movement. Very few articles reported household income and ethnicity/race. Key themes investigated by this body of literature were breast biomechanics, breast pain, and physical activity levels. Meta-analyses revealed that women and girls with larger breasts reported more exercise-induced breast pain and experienced greater breast mediolateral velocity and anterior/poster acceleration during physical activity than women and girls with smaller breasts. Meta-analyses revealed that sports bras were associated with less breast pain during physical activity than standard bras.

CONCLUSION: This body of literature largely studied a homogenous population, and there is a need for greater inclusion of women and girls who identify with under-represented minorities, with disability, who are post-menopausal, and who are overweight. Factors such as ethnicity/race and household income, were underreported in the literature, and studies of movement patterns beyond running are needed. Mediolateral breast velocity and anterior/posterior breast acceleration may be contributors to breast pain in women and girls with larger breasts, thus these biomechanical outcomes are of particular interest for future research and for consideration in sports bra optimization.



INTRODUCTION

Before 1986, women* were largely excluded from clinical trials and research out of concern for impacts on fertility.¹ This exclusion expanded to areas of research that do not put fertility at risk, such as physical activity, and to this day still negatively impacts our understanding of physical activity in females.² Indeed, women often have lower levels of physical activity and higher rates of obesity as compared to men.³-6 This activity gap is even greater for women of color.7 Although the drivers of these disparities are multifactorial, breast pain and motion are one of the significant barriers to women's participation in physical activity and sport.8

In 1978, Haycock et al. first documented how breast motion impacts running and jumping amongst females.9 Their survey found that 31% of female athletes experienced breast soreness following exercise. Indeed, the breasts can move up to 15 cm during vigorous physical activity, and excessive breast motion is associated with breast discomfort, affecting as many as 56% of females. 10-13 In fact, one study suggested as little as 2 cm of breast motion during exercise is associated with discomfort.14 Unfortunately, this breast pain often deters women and girls from exercising and participating in sports. In a survey of 249 women in the United Kingdom, 24% reported reducing their physical activity levels to relieve breast pain and 33% reported not meeting physical activity guidelines as a result.8 Thus, despite the fact that knowledge of breast impacts on physical activity have existed for over 40 years, there are still breast support obstacles for women and girls during physical activity.

In examining components contributing to exercise-induced breast pain, breast size is an important factor. Specifically, females with larger breasts exhibit greater breast motion and report more breast pain during running and jumping activities. 11,13,15 In a survey of over 2,000 adolescent girls in the United Kingdom, 63% of larger-breasted school girls reported their breasts had a negative impact on their participation in sports and exercise, citing excess breast motion as their greatest concern. 16 Similarly, a survey of 438 females in the

United States demonstrated that females with larger breasts were significantly more likely to report that concerns of breast support prevented them from participating in physical activity.¹⁷ The impact of large breasts disproportionately affects women who are overweight or obese, and increasing breast size has been found to correlate with a decreasing physical activity cycle, which further contributes to and worsens rates of obesity.^{14,18,19,20,21} Therefore, the impact of breasts and bras on physical activity disproportionately affects women and girls with larger breasts and other specific populations.

To try to mitigate some of the breast-motion related barriers to physical activity, in 1977, the first activity-specific bra, known as the "jogbra", was designed.²² ²³ Decades later, sports bras have grown into a multibillion-dollar industry and are an essential piece of sporting equipment for many women and girls to be comfortable while being active and participating in sports.^{24,25} Although in research studies sports bras appear to reduce breast pain and movement during physical activity, breast pain continues to be a reported barrier to physical activity, even as recent as 2023. 10,12,26,27 Thus, there is a need to synthesize the literature aimed at understanding how breasts and bras impact physical activity to identify potential explanations as to why breast-related barriers to physical activity still exist.

With this background in mind, the purpose of this systematic review was to describe how bras and breasts impact physical activity-related outcomes in girls and women. To be as all-encompassing as possible, we used the World Health Organizations definition of physical activity, defined as all movement including but not limited to exercise, sports, play, and walking.28 We hypothesized that girls and women with larger breasts would (i) have more pain and discomfort with physical activity, (ii) have more barriers to physical activity, and (iii) be less physically active than women and girls with smaller breasts. We also hypothesized that improper support of breasts during physical activity will lead to (i) improper biomechanics, (ii) more pain during physical activity, and (iii) more injuries than proper breast support.

on the literature that is available, thus, we use the terms "woman" and "girl" to describe individuals who identify as a woman or girl and has breasts.

^{*} While in this review we focus on the impacts of bras and breasts on physical activity in women and girls, we acknowledge that not all women and girls have breasts and that some individuals with breasts do not identify as a woman or girl. Our focus here is largely limited based



METHODS

Systematic Review

Our systematic review was registered on Open Science Framework (OSF) Registries a priori under the title "The impact of breasts and bras on females' ability to move: a scoping review". A literature search was performed under the guidance of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement and protocol for systematic reviews (Appendix S1), the Meta-Analysis of Observational Studies in Epidemiology (MOOSE) checklist (Appendix S2), and the Cochrane Handbook for Systematic Reviews of Interventions.²⁹⁻³² Studies were not excluded based on the presence of a control group (e.g., case study), age of participants, or year of publications. Studies were excluded if the paper was not published in a peer reviewed journal, not written in English, did not contain human subjects, was not an original research article, studied males only, or focused on breastfeeding/post-partum or breast cancer.

We performed electronic searches of PubMed, Physiotherapy Evidence Database (PEDro). Cumulative Index to Nursing and Allied Health Literature (CINAHL), and Cochrane Central Register of Controlled Trials on March 24, 2023, with search terms outlined in Table S1. Title, abstract, and full-text screening was performed by independent reviewers (XX, Disagreements were resolved by an independent third reviewer (XX). All screening steps were completed in pre-designed spreadsheets.

Two reviewers extracted study information from the included publications. Specifically, we documented the variables listed under "Study/Participant Demographics Cataloged" in Table S2. If these data were not reported in an included study, we noted so. We then further

categorized studies based on intervention type and outcome variables (see Table S2 for specific categories considered).

Meta-analyses

We performed meta-analyses on overlapping outcomes that were reported amongst studies. Specifically, this included evaluating the effect of (i) breast size on breast pain, physical activity level, and breast biomechanics, (ii) bra type on breast pain, and (iii) breast reduction on physical activity level. Breast size was categorized into small breasts (A-C cup) and large breasts (D cup or larger) since this was the most commonly utilized reductionist approach amongst included studies, and bra type was categorized into standard bra and sports bra since these were the two most commonly investigated bra types. Standardized mean differences (SMD) and pooled standard deviations (SD_{pool}) of outcome measures were calculated using the DerSimonian-Laird method.³² All metaanalyses were performed using SPSS Statistics for Windows (version 28.0, IBM Corp., Armonk, NY, USA).

RESULTS

Literature aimed at studying the impact of breasts/bras on physical activity is largely focused on a homogenous study population

From our systematic literature search, we identified 323 articles studying the effect of breasts and/or bras on physical activity. Following title, abstract, and full text screening, 77 articles were eligible for inclusion in our review (Figure 1).8,10-14,16-18,20,21,26,33-97 Reasons for exclusion at the full text level are listed in Table S3, with the most frequent reason for exclusion being the article was not an original research study.






Figure 1. Systematic review workflow. Systematic search of databases, abstract, title, and full text screening resulted in 77 unique articles included in this review

Details on exclusion reasons at the full text level and included article details are available in Tables S3 and S4

Detailed study information was extracted from the included articles and is listed in Table S3. Approximately half of the articles focused on women ages 20-29 (n = 32, Figure 2A), and the majority of articles did not report the ethnicity/race of their participants (n = 73, Figure 2B). Of those that did, most participants were white (70% or more, Table S4). Additionally, only one study reported the household income of participants.¹⁷ Published studies largely took place in Australia, the United Kingdom, or the United States (n = 71, Figure 2C), and included studies mainly focused on nonathletes (n = 62, Figure 2D). Of those that did include athletes, runners were the most commonly studied population (n = 9/17, (two studies included both athletes and non-athletes) Figure 2D). Some studies reported participants' weight (n = 52) or body mass index (BMI) (n = 18), with the mean weight being 45-65 kg and the most investigated BMI being 18.5-29.9; however, a fifth of studies did not report the weight of their participants (n = 14, Figure 2E) and the majority (n = 59) did not report BMI (Figure 2F). There was a wide range of breast sizes represented amongst participants in included studies, with the most common cup size being D (Figure 2G). Most studies were Level III or IV evidence (n = 58, Figure 2H) and focused on biomechanical outcomes (n = 43, Figure 2I). Additionally, the average impact factor of journals from included articles was 4.0 ± 3.4, and 6% of studies were published in female-focused journals (e.g., The Breast Journal). Encouragingly, the rate of publication of literature aimed at studying how breasts and bras impact physical activity is increasing exponentially across time (Figure 2J). In examining the specific movement under study, most articles focused on running (n = 41, Figure 2K; Table S4 list movement under study on a per article



basis). Many studies' independent variable was bra type (n = 37), while 23 studies focused on breast size, and 24 studies were descriptive in nature (Figure 2L). The most common outcome variables studied in this body of literature were related to breast biomechanics (n = 31) and breast pain (n = 31, Figure 2M).

Meta-analyses revealed that breasts and bras impact physical activity level and breast pain during physical activity

Meta-analyses revealed that women with larger breasts reported more breast pain with physical activity than women with smaller breasts (Figure 3A). Despite this, however, there was no detectable effect of breast size on physical activity level (Figure 3B). Interestingly, sports bras were reported to cause less breast pain during physical activity than standard bras (Figure 3C), and women reported more physical activity post-breast reduction than pre-breast reduction (Figure 3D).

Women with larger breasts experienced greater mediolateral velocity and anterior/posterior acceleration of their breasts during physical activity than women with smaller breasts

When examining the effect of breast size on breast displacement, we found no effect in the mediolateral, or anterior/posterior vertical, directions (Figure 4A-4C). While there was no effect of breast size on vertical and anterior/posterior velocity (Figure 4D, 4F), women and girls with larger breasts experienced greater mediolateral velocity than women and girls with smaller breasts (Figure 4E). Conversely, while there was no effect of breast size on vertical and mediolateral acceleration (Figure 4G-4H), women and girls with larger breasts experienced greater anterior/posterior acceleration than women and girls with smaller breasts (Figure 4I).

DISCUSSION

This systematic review provides a comprehensive summary of the existing literature aimed at understanding the impact of breasts and bras on physical activity amongst women and girls. While there is a large body of literature reviewing nonbreast related physical activity outcomes amongst females, to our knowledge, there are only three published reviews focused specifically on breasts

and physical activity outcomes.⁹⁸⁻¹⁰⁹ Two of these reviews specifically examine methodology for studying breast biomechanics, while the most recently published review analyzed how bra fit relates to sports participation.¹¹⁰⁻¹¹² Here, we build on this body of literature by more broadly examining how breasts and bras impact physical activity and performing meta-analyses on how these factors impact breast pain, physical activity levels, and breast biomechanics.

Encouragingly, the body of literature aimed at studying the impact of breasts and bras on physical activity is growing exponentially and is being published in journals geared toward the entire exercise/sports/rehabilitation community. Indeed, we found that the number of publications on this topic increased more than 3-fold from 2003-2012 to 2013-2022. Additionally, we found that 94% of studies were published in general sports/exercise/rehabilitation journals, while only 6% of studies were published in female-specific journals. These findings are of particular interest, given the consistently reported lack of studies on female athletes relative to male counterparts and more generally challenges with publishing studies that include women compared to studies that include men.^{113,114}

The literature revealed homogenous subject populations, with the majority of articles examining participants who were 20-29 years old, identified as white, lived in Western countries, were nonathletes, and weighed between 45-65 Encouragingly, however, many studies included a wide range of breast sizes (Figure 2F). In future studies, greater attention should be paid to the inclusion of older women, women with disabilities, underrepresented minorities, and overweight or obese women. Interestingly, race and ethnicity, along with household income and other social determinants of health, were not reported by many studies. When studies did report race, the vast majority of participants identified as white. As of 2023, all but six studies were conducted in predominantly white countries including Australia, the United Kingdom, and the United States. As sports bra-wearing habits and perceptions of the breasts likely vary in different communities of women, the results gathered in this review may not be generalizable to other populations, and greater attention to including diverse populations is critical for future research.






Figure 2. Descriptors of publications aimed at studying the effects of breasts and bras on physical activity in women and girls. A) Ages included, B) Race/Ethnicity, C) Location of study, D) Athlete status, E) Weight of included participants, F) Body mass index (BMI) of participants, G) Breast size of participants, H) Level of evidence, I) Study Type (i.e., what types of data were collected/what was the format of the study), J) Year of publication, K) Movement type under study, L) Intervention type or independent variable of the study, M) Outcome variables or dependent variables in study






Figure 3. The effects of breasts and bras on breast pain during physical activity and physical activity level

Meta-analyses were performed to assess the following relationships: (A) the effect of breast size on breast pain during physical activity, (B) the effect of breast size on physical activity level, (C) the effect of bra type on breast pain during physical activity, and (D) the effect of breast reduction on physical activity level.

Few studies were specifically focused on athletes. When they did, most articles analyzing athletes included only runners, and the most common movement-related intervention was running. Only two of the included studies examined breast outcomes amongst athletes in contact sports. In a survey of 504 elite female athletes representing 46 different participants in contact sports reported a significantly greater number of breast injuries (47.3%) than participants in non-contact sports (33.7%).82 Another study reported that amongst collegiate-level female basketball and soccer players, 48.8% and 46.7%, respectively, had experienced a breast injury while playing their sport.80 While the data is limited, these studies suggest that athletes, especially those in contact sports, experience high rates of breast injuries. Future research on this topic could focus on understanding the causes and types of breast injuries experienced by contact sport players.

In addition to breast injuries amongst athletes, exercise-induced breast pain presents a significant burden to physical activity. In a survey of elite female athletes in Australia, 44% reported experiencing exercise-induced breast pain, with 32% reporting this pain negatively affected performance in their sport. 115 Of the few studies that have examined other modalities of physical activity, results suggest breast biomechanics perceptions of breast impacts on physical activity vary considerably between walking, running, and jumping.86,110 Thus, our understanding of the impact of breasts and bras on athletes and in sports/movements outside of running, especially in contact sports, is limited and is a needed focus area of future research.

Of the studies included in our review, five specifically studied bra fit, though not in a manner we could specifically compile into a meta-analysis. When asking non-athlete young women with C-E breast size about their experience with bra fitting, 87% reported having never been fitted professionally.⁶³ In addition, 83% of female athletes



at the Olympics and Paralympics wore improperly fitted bras.94 Another study reported that bra fit discomfort increases during running when a standard sports bra is used, and Wakefield-Scurr et al. reported that sports bra fit changes after 25 wash/dry cycles. 47,95 Given this current literature, there is an essential need for more access to bra fitting and educational initiatives, for athletes and non-athletes alike, on appropriate bra fit as well as descriptive studies geared at understanding how much inappropriate bra fit, access to sports bra, and poor sports bra design are barriers for physical activity.

Interestingly, our meta-analyses did not detect an effect of breast size on physical activity level. This lack of effect may be due to missing confounders, such as information on bra fit or type. We did, however, identify that larger breasts were associated with both more breast pain and altered breast biomechanics. Specifically, we found women and girls with larger breasts presented with more mediolateral velocity and anterior/posterior acceleration of their breasts compared to women with smaller breasts. This correlation between breast biomechanics and pain puts forth the hypothesis that increased mediolateral velocity and anterior/poster acceleration of the breasts may contribute to breast pain. However, similar to the meta-analysis on breast size and physical activity level, these data are also limited by incomplete reporting of bra fit and type, which may affect the conclusions drawn by this dataset.

The effect of breast size on breast biomechanics during physical activity




Figure 4. The effects of breast size on breast biomechanics during physical activity. Meta-analyses were performed to assess the effect of breast size on breast: A) vertical displacement, B) mediolateral displacement, C) anterior/posterior displacement, D) vertical velocity, E) mediolateral velocity, F) anterior/posterior velocity, G) vertical acceleration, H) mediolateral acceleration, and I) anterior/posterior acceleration

Our meta-analyses indicate that a standard | during physical activity in comparison to a regular sports bra is sufficient at reducing breast pain | bra. While this does not negate the importance of



continued improvement on sports bra design, it does raise into question why breast pain continues to be such a prominent problem for females, both athletes and non-athletes, during physical activity.8 Indeed, many studies have suggested that wearing a properly fitted sports bra can reduce breast movement by 59% and decrease pain in 85% of cases. 10,116 As aforementioned, there is evidence that access to bra fitting and/or education on the importance of bra fitting is missing amongst women included within these studies. There is also anecdotal evidence that access to sports bras, appropriately fitted or not, is a central barrier to particular communities of women to physical activity, and there are many organizations aimed at increasing sports bra accessibility (for example, The Sports Bra Project and Bras for Girls). 117-122 However, this is a poorly reported and studied area of the literature, and only one study in our review addressed this head on.¹⁷ Therefore, understanding how access to sports bras impacts breast pain during physical activity and improving said access is an important area of continued research and outreach efforts.

Although this study adds to a growing body of literature, it does exist with limitations. As described previously, the study populations and movement patterns examined in the literature are largely homogenous, thus generalizability to other populations and movements limited. Additionally, given the limited types of physical activity studied, we were not able to stratify data by activity type. The included studies also sparsely reported baseline activity level, BMI, bra type*, bra fit*, accessibility of sports bras, and other health risk factors that may have confounded results; thus, we were not able to adequately control for these features (*unless these were the specific independent/dependent variables). Another limitation of our analyses is that these studies were published over a wide range of time (1973-2023), and bra design and support has changed over this time period in a way that we have not controlled for here. As mentioned at the onset of our study, these analyses exclude people who have breasts but do not identify as women/girls, and there is a need for specific research aimed at targeting this population. Our meta-analyses were limited by a small sample size and presented with high heterogeneity ($I^2 > 0.9$ for all), thus the lack of effects may be due to underpowered analysis or confounding factors rather than a true lack of effect. Additionally, we did not perform risk of bias or rigor and reproducibility analyses.

CONCLUSION

This systematic review and meta-analysis studying the impact of breasts and bras on physical activity in women and girls. We found that the literature is biased towards a homogenous population, and greater inclusion of underrepresented minorities, post-menopausal, and overweight women will increase the generalizability of studies. In addition, there is a need for this body of literature to better report confounding factors, such as race/ethnicity and household income, and study other types of movement beyond running. Our meta-analyses revealed that larger breasts were associated with both more breast pain, more mediolateral breast velocity, and more anterior/posterior breast acceleration. Thus, prospective studies should focus on these biomechanical outcomes in sports bra design to further determine if these altered mechanics have a causal relationship with breast pain.

Acknowledgments

This study was conducted as a part of the Female Athlete Research Interest Group of the Pediatric Research in Sports Medicine Society.

Conflict of Interest Statement

This study was financially supported by the following grant: T32AG021885-19 for GG. The authors declare no other conflicts of interest with the contents of this study.

Corresponding Author

Gabrielle Gilmer, BChE University of Pittsburgh School of Medicine M247B Scaife Hall 3550 Terrace Street Pittsburg, Pennsylvania 15261

Email: ggg15@pitt.edu
Telephone: 205-789-9607



REFERENCES

- Merkatz RB. Inclusion of women in clinical trials: a historical overview of scientific, ethical, and legal issues. J Obstet Gynecol Neonatal Nurs. 1998;27(1):78-84. Epub 1998/02/25. doi: 10.1111/j.1552-6909.1998.tb02594.x. PubMed PMID: 9475131.
- McCarthy CR. Historical background of clinical trials involving women and minorities. Acad Med. 1994;69(9):695-8. Epub 1994/09/01. doi: 10.1097/00001888-199409000-00002. PubMed PMID: 8074757.
- 3. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. Med Sci Sports Exerc. 2008;40(1):181-8. Epub 2007/12/20. doi: 10.1249/mss.0b013e31815a51b3. PubMed PMID: 18091006.
- Palmer MK, Toth PP. Trends in Lipids, Obesity, Metabolic Syndrome, and Diabetes Mellitus in the United States: An NHANES Analysis (2003-2004 to 2013-2014). Obesity (Silver Spring). 2019;27(2):309-14. Epub 2019/01/25. doi: 10.1002/oby.22370. PubMed PMID: 30677260.
- Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of Obesity and Severe Obesity Among Adults: United States, 2017-2018. NCHS Data Brief. 2020(360):1-8. Epub 2020/06/04. PubMed PMID: 32487284.
- Ogden CL, Fryar CD, Martin CB, Freedman DS, Carroll MD, Gu Q, et al. Trends in Obesity Prevalence by Race and Hispanic Origin-1999-2000 to 2017-2018. JAMA. 2020;324(12):1208-10. Epub 2020/08/29. doi: 10.1001/jama.2020.14590. PubMed PMID: 32857101; PubMed Central PMCID: PMCPMC7455882.
- 7. Saffer H, Dave D, Grossman M, Leung LA. Racial, Ethnic, and Gender Differences in Physical Activity. J Hum Cap. 2013;7(4):378-410. Epub 2013/01/01. doi: 10.1086/671200. PubMed PMID: 25632311; PubMed Central PMCID: PMCPMC4306423.
- 8. Burnett E, White J, Scurr J. The Influence of the Breast on Physical Activity Participation in Females. J Phys Act Health. 2015;12(4):588-94. Epub 2014/06/07. doi: 10.1123/jpah.2013-0236. PubMed PMID: 24905581.
- 9. Haycock CE SGaGJ. The female athlete does her anatomy pose problems? Proceedings of the American Medical Association 19th Conference on the Medical Aspects of Sports; Monroe, WI: AMA Press; 1978. p. 1-8.
- Scurr JC, White JL, Hedger W. Supported and unsupported breast displacement in three dimensions across treadmill activity levels. J Sports Sci. 2011;29(1):55-61. Epub 2010/11/16. doi: 10.1080/02640414.2010.521944. PubMed PMID: 21077006.
- 11. Lorentzen D, Lawson L. Selected Sports Bras: A Biomechanical Analysis of Breast Motion While

- Jogging. Phys Sportsmed. 1987;15(5):128-39. Epub 1987/05/01. doi: 10.1080/00913847.1987.11709355. PubMed PMID: 27463876.
- 12. Mason BR, Page KA, Fallon K. An analysis of movement and discomfort of the female breast during exercise and the effects of breast support in three cases. J Sci Med Sport. 1999;2(2):134-44. Epub 1999/09/07. doi: 10.1016/s1440-2440(99)80193-5. PubMed PMID: 10476977.
- 13. Bridgman C, Scurr J, White J, Hedger W, Galbraith H. Three-dimensional kinematics of the breast during a two-step star jump. J Appl Biomech. 2010;26(4):465-72. Epub 2011/01/20. doi: 10.1123/jab.26.4.465. PubMed PMID: 21245506.
- 14. Coltman CE, Steele JR, McGhee DE. Does breast size affect how women participate in physical activity? J Sci Med Sport. 2019;22(3):324-9. Epub 2018/09/25. doi: 10.1016/j.jsams.2018.09.226. PubMed PMID: 30244978.
- Gehlsen G, Albohm M. Evaluation of Sports Bras. Phys Sportsmed. 1980;8(10):88-97. Epub 1980/10/01. doi: 10.1080/00913847.1980.11948653. PubMed PMID: 29261415.
- Scurr J, Brown N, Smith J, Brasher A, Risius D, Marczyk A. The Influence of the Breast on Sport and Exercise Participation in School Girls in the United Kingdom. J Adolesc Health. 2016;58(2):167-73. Epub 2016/01/24. doi: 10.1016/j.jadohealth.2015.10.005. PubMed PMID: 26802991.
- 17. Rizzone KH, Edison B, Coleman N, Carter C, Ichesco I, Cassidy P, et al. Sports Bra Preferences by Age and Impact of Breast Size on Physical Activity among American Females. Int J Environ Res Public Health. 2021;18(23). Epub 2021/12/11. doi: 10.3390/ijerph182312732. PubMed PMID: 34886465; PubMed Central PMCID: PMCPMC8657319.
- Steele JR, Coltman CE, McGhee DE. Effects of obesity on breast size, thoracic spine structure and function, upper torso musculoskeletal pain and physical activity in women. J Sport Health Sci. 2020;9(2):140-8. Epub 2020/02/27. doi: 10.1016/j.jshs.2019.05.003. PubMed PMID: 32099722; PubMed Central PMCID: PMCPMC7031809.
- Lubian Lopez DM, Butron Hinojo CA, Castillo Lara M, Sanchez-Prieto M, Sanchez-Borrego R, Mendoza Ladron de Guevara N, et al. Relationship of breast volume, obesity and central obesity with different prognostic factors of breast cancer. Sci Rep. 2021;11(1):1872. Epub 2021/01/23. doi: 10.1038/s41598-021-81436-9. PubMed PMID: 33479372; PubMed Central PMCID: PMCPMC7820412.
- Oon IH, Mara JK, Steele JR, McGhee DE, Lewis V, Coltman CE. Women with larger breasts are less satisfied with their breasts: Implications for quality of life and physical activity participation. Womens Health (Lond). 2022;18:17455057221109394. Epub 2022/07/09. doi: 10.1177/17455057221109394.



- PubMed PMID: 35801682; PubMed Central PMCID: PMCPMC9274428.
- Spencer L, Fary R, McKenna L, Jacques A, Lalor J, Briffa K. The relationship between breast size and aspects of health and psychological wellbeing in mature-aged women. Womens Health (Lond). 2020;16:1745506520918335. Epub 2020/05/19. doi: 10.1177/1745506520918335. PubMed PMID: 32419664; PubMed Central PMCID: PMCPMC7235664.
- 22. Schuster K. Equipment Update: Jogging Bras Hit the Streets. Phys Sportsmed. 1979;7(4):125-8. Epub 1979/04/01. doi: 10.1080/00913847.1979.11710846. PubMed PMID: 27421500.
- 23. Keyes A. How the First Sports Bra Got Its Stabilizing Start. Smithsonian Magazine. 2020.
- 24. Shah R. Sports Bras Market is expected to reach USD 103.56 billion by 2029, at a CAGR of 11.4%: GreyViews: GlobalNewWire; 2022 [cited 2023 July 28]. Available from: https://www.globenewswire.com/news-release/2022/12/02/2566671/0/en/Sports-Bras-Market-is-expected-to-reach-USD-103-56-billion-by-2029-at-a-CAGR-of-11-4-GreyViews.html.
- 25. Jeroen Van Heghe CS. Global Bra Market Research Report - Industry Analysis, Size, Share, Growth, Trends and Forecast 2023 to 2030. Value Market Research, Global Information. 2023.
- Gibson TM, Balendra N, Ustinova KI, Langenderfer JE. Reductions in Kinematics from Brassieres with Varying Breast Support. Int J Exerc Sci. 2019;12(1):402-11. Epub 2019/03/23. PubMed PMID: 30899340; PubMed Central PMCID: PMCPMC6413842.
- Baxter CR, Crittenden TA, Dean NR. Self-reported breast size, exercise habits and BREAST-Q data - an international cross-sectional study of community runners. JPRAS Open. 2023;37:92-101. Epub 2023/07/17. doi: 10.1016/j.jpra.2023.06.013. PubMed PMID: 37457991; PubMed Central PMCID: PMCPMC10339238.
- 28. Organization WH. Physical activity 2023 [cited 2023]. Available from: https://www.who.int/health-topics/physical-activity#tab=tab_1.
- 29. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Ann Intern Med. 2009;151(4):264-9, W64. Epub 2009/07/23. doi: 10.7326/0003-4819-151-4-200908180-00135. PubMed PMID: 19622511.
- Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015;350:g7647. Epub 2015/01/04. doi: 10.1136/bmj.g7647. PubMed PMID: 25555855.
- 31. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of

- observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA. 2000;283(15):2008-12. Epub 2000/05/02. doi: 10.1001/jama.283.15.2008. PubMed PMID: 10789670.
- 32. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Cochrane Handbook for Systematic Reviews of Interventions. 6.3 ed2022.
- Hunter LY, Torgan C. The Bra Controversy: Are Sports Bras a Necessity? Phys Sportsmed. 1982;10(11):75-6. Epub 1982/11/01. doi: 10.1080/00913847.1982.11947369. PubMed PMID: 29291314.
- Glatt BS, Sarwer DB, O'Hara DE, Hamori C, Bucky LP, LaRossa D. A retrospective study of changes in physical symptoms and body image after reduction mammaplasty. Plast Reconstr Surg. 1999;103(1):76-82; discussion 3-5. Epub 1999/01/23. doi: 10.1097/00006534-199901000-00013. PubMed PMID: 9915166.
- Blomqvist L, Eriksson A, Brandberg Y. Reduction mammaplasty provides long-term improvement in health status and quality of life. Plast Reconstr Surg. 2000;106(5):991-7. Epub 2000/10/20. doi: 10.1097/00006534-200010000-00005. PubMed PMID: 11039369.
- Mizgala CL, MacKenzie KM. Breast reduction outcome study. Ann Plast Surg. 2000;44(2):125-33; discussion 33-4. Epub 2000/03/01. doi: 10.1097/00000637-200044020-00001. PubMed PMID: 10696037.
- 37. Netscher DT, Meade RA, Goodman CM, Brehm BJ, Friedman JD, Thornby J. Physical and psychosocial symptoms among 88 volunteer subjects compared with patients seeking plastic surgery procedures to the breast. Plast Reconstr Surg. 2000;105(7):2366-73. Epub 2000/06/09. doi: 10.1097/00006534-200006000-00009. PubMed PMID: 10845288.
- 38. Sood R, Mount DL, Coleman JJ, 3rd, Ranieri J, Sauter S, Mathur P, et al. Effects of reduction mammaplasty on pulmonary function and symptoms of macromastia. Plast Reconstr Surg. 2003;111(2):688-94. Epub 2003/02/01. doi: 10.1097/01.PRS.0000041395.02699.B7. PubMed PMID: 12560689.
- 39. Bowles KA, Steele JR, Chaunchaiyakul R. Do current sports brassiere designs impede respiratory function? Med Sci Sports Exerc. 2005;37(9):1633-40. Epub 2005/09/24. doi: 10.1249/01.mss.0000177590.75686.28. PubMed PMID: 16177619.
- Williams PT. Nonlinear relationships between weekly walking distance and adiposity in 27,596 women. Med Sci Sports Exerc. 2005;37(11):1893-901. Epub 2005/11/16. doi: 10.1249/01.mss.0000175860.51204.85. PubMed PMID: 16286858; PubMed Central PMCID: PMCPMC2813189.



- 41. McGhee DE, Power BM, Steele JR. Does deep water running reduce exercise-induced breast discomfort? Br J Sports Med. 2007;41(12):879-83; discussion 83. Epub 2007/05/31. doi: 10.1136/bjsm.2007.036251. PubMed PMID: 17535854; PubMed Central PMCID: PMCPMC2658982.
- 42. Spector JA, Karp NS. Reduction mammaplasty: a significant improvement at any size. Plast Reconstr Surg. 2007;120(4):845-50. Epub 2007/09/07. doi: 10.1097/01.prs.0000277660.49802.76. PubMed PMID: 17805110.
- Bowles KA, Steele JR, Munro B. What are the breast support choices of Australian women during physical activity? Br J Sports Med. 2008;42(8):670-3. Epub 2008/06/05. doi: 10.1136/bjsm.2008.046219. PubMed PMID: 18523041.
- 44. White JL, Scurr JC, Smith NA. The effect of breast support on kinetics during overground running performance. Ergonomics. 2009;52(4):492-8. Epub 2009/04/30. doi: 10.1080/00140130802707907. PubMed PMID: 19401901.
- 45. Scurr J, White J, Hedger W. Breast displacement in three dimensions during the walking and running gait cycles. J Appl Biomech. 2009;25(4):322-9. Epub 2010/01/26. doi: 10.1123/jab.25.4.322. PubMed PMID: 20095453.
- 46. McGhee DE, Steele JR, Munro BJ. Education improves bra knowledge and fit, and level of breast support in adolescent female athletes: a cluster-randomised trial. J Physiother. 2010;56(1):19-24. Epub 2010/05/27. doi: 10.1016/s1836-9553(10)70050-3. PubMed PMID: 20500133.
- McGhee DE, Steele JR. Breast elevation and compression decrease exercise-induced breast discomfort. Med Sci Sports Exerc. 2010;42(7):1333-8. Epub 2009/12/19. doi: 10.1249/MSS.0b013e3181ca7fd8. PubMed PMID: 20019639.
- 48. Scurr JC, White JL, Hedger W. The effect of breast support on the kinematics of the breast during the running gait cycle. J Sports Sci. 2010;28(10):1103-9. Epub 2010/08/06. doi: 10.1080/02640414.2010.497542. PubMed PMID: 20686995.
- Singh KA, Pinell XA, Losken A. Is reduction mammaplasty a stimulus for weight loss and improved quality of life? Ann Plast Surg. 2010;64(5):585-7. Epub 2010/04/01. doi: 10.1097/SAP.0b013e3181c5a841. PubMed PMID: 20354428.
- 50. White J, Scurr J, Hedger W. A comparison of three-dimensional breast displacement and breast comfort during overground and treadmill running. J Appl Biomech. 2011;27(1):47-53. Epub 2011/04/01. doi: 10.1123/jab.27.1.47. PubMed PMID: 21451182.
- 51. Bowles KA, Steele JR, Munro BJ. Features of sports bras that deter their use by Australian women. J Sci Med Sport. 2012;15(3):195-200. Epub 2011/12/23.

- doi: 10.1016/j.jsams.2011.11.248. PubMed PMID: 22188847.
- 52. Wood LE, White J, Milligan A, Ayres B, Hedger W, Scurr J. Predictors of three-dimensional breast kinematics during bare-breasted running. Med Sci Sports Exerc. 2012;44(7):1351-7. Epub 2012/06/19. doi: 10.1249/MSS.0b013e31824bd62c. PubMed PMID: 22705551.
- 53. McGhee DE, Steele JR, Zealey WJ, Takacs GJ. Brabreast forces generated in women with large breasts while standing and during treadmill running: Implications for sports bra design. Appl Ergon. 2013;44(1):112-8. Epub 2012/06/30. doi: 10.1016/j.apergo.2012.05.006. PubMed PMID: 22742876.
- 54. Bowles KA, Steele JR. Effects of strap cushions and strap orientation on comfort and sports bra performance. Med Sci Sports Exerc. 2013;45(6):1113-9. Epub 2013/01/01. doi: 10.1249/MSS.0b013e3182808a21. PubMed PMID: 23274596.
- 55. Ayres B, White J, Hedger W, Scurr J. Female upper body and breast skin temperature and thermal comfort following exercise. Ergonomics. 2013;56(7):1194-202. Epub 2013/05/09. doi: 10.1080/00140139.2013.789554. PubMed PMID: 23651438.
- Brown N, White J, Brasher A, Scurr J. The experience of breast pain (mastalgia) in female runners of the 2012 London Marathon and its effect on exercise behaviour. Br J Sports Med. 2014;48(4):320-5. Epub 2013/04/23. doi: 10.1136/bjsports-2013-092175. PubMed PMID: 23603819.
- 57. Milligan A, Mills C, Scurr J. The effect of breast support on upper body muscle activity during 5 km treadmill running. Hum Mov Sci. 2014;38:74-83. Epub 2014/09/26. doi: 10.1016/j.humov.2014.06.001. PubMed PMID: 25255202.
- 58. Milligan A, Mills C, Scurr J. Within-participant variance in multiplanar breast kinematics during 5 km treadmill running. J Appl Biomech. 2014;30(2):244-9. Epub 2013/10/23. doi: 10.1123/jab.2013-0121. PubMed PMID: 24146061.
- 59. Mills C, Loveridge A, Milligan A, Risius D, Scurr J. Can axes conventions of the trunk reference frame influence breast displacement calculation during running? J Biomech. 2014;47(2):575-8. Epub 2013/12/18. doi: 10.1016/j.jbiomech.2013.11.041. PubMed PMID: 24342499.
- 60. Mills C, Loveridge A, Milligan A, Risius D, Scurr J. Is torso soft tissue motion really an artefact within breast biomechanics research? J Biomech. 2014;47(11):2606-10. Epub 2014/06/24. doi: 10.1016/j.jbiomech.2014.05.023. PubMed PMID: 24953261.
- 61. Brown N, White J, Brasher A, Scurr J. An investigation into breast support and sports bra use



- in female runners of the 2012 London Marathon. J Sports Sci. 2014;32(9):801-9. Epub 2014/02/01. doi: 10.1080/02640414.2013.844348. PubMed PMID: 24479370.
- 62. White J, Mills C, Ball N, Scurr J. The effect of breast support and breast pain on upper-extremity kinematics during running: implications for females with large breasts. J Sports Sci. 2015;33(19):2043-50. Epub 2015/03/21. doi: 10.1080/02640414.2015.1026378. PubMed PMID: 25793936.
- 63. Coltman CE, McGhee DE, Steele JR. Bra strap orientations and designs to minimise bra strap discomfort and pressure during sport and exercise in women with large breasts. Sports Med Open. 2015;1(1):21. Epub 2015/08/19. doi: 10.1186/s40798-015-0014-z. PubMed PMID: 26284162; PubMed Central PMCID: PMCPMC5005736.
- 64. Risius D, Milligan A, Mills C, Scurr J. Multiplanar breast kinematics during different exercise modalities. Eur J Sport Sci. 2015;15(2):111-7. Epub 2014/06/20. doi: 10.1080/17461391.2014.928914. PubMed PMID: 24942053.
- 65. Milligan A, Mills C, Corbett J, Scurr J. The influence of breast support on torso, pelvis and arm kinematics during a five kilometer treadmill run. Hum Mov Sci. 2015;42:246-60. Epub 2015/06/17. doi: 10.1016/j.humov.2015.05.008. PubMed PMID: 26079773.
- Mills C, Ayres B, Scurr J. Breast Support Garments are Ineffective at Reducing Breast Motion During an Aqua Aerobics Jumping Exercise. J Hum Kinet. 2015;46:49-58. Epub 2015/08/05. doi: 10.1515/hukin-2015-0033. PubMed PMID: 26240648; PubMed Central PMCID: PMCPMC4519221.
- Mills C, Risius D, Scurr J. Breast motion asymmetry during running. J Sports Sci. 2015;33(7):746-53. Epub 2014/10/31. doi: 10.1080/02640414.2014.962575. PubMed PMID: 25356791.
- 68. Milligan A, Mills C, Corbett J, Scurr J. Magnitude of multiplanar breast kinematics differs depending upon run distance. J Sports Sci. 2015;33(19):2025-34. Epub 2015/04/02. doi: 10.1080/02640414.2015.1026376. PubMed PMID: 25827174.
- 69. Mills C, Lomax M, Ayres B, Scurr J. The movement of the trunk and breast during front crawl and breaststroke swimming. J Sports Sci. 2015;33(4):427-36. Epub 2014/09/06. doi: 10.1080/02640414.2014.946951. PubMed PMID: 25189996.
- 70. Schinkel-Ivy A, Drake JD. Breast size impacts spine motion and postural muscle activation. J Back Musculoskelet Rehabil. 2016;29(4):741-8. Epub 2016/03/12. doi: 10.3233/BMR-160680. PubMed PMID: 26966826.
- 71. Nolte K, Burgoyne S, Nolte H, J VDM, Fletcher L. The effectiveness of a range of sports bras in reducing breast displacement during treadmill

- running and two-step star jumping. J Sports Med Phys Fitness. 2016;56(11):1311-7. Epub 2016/01/15. PubMed PMID: 26765495.
- 72. Brown N, Scurr J. Do women with smaller breasts perform better in long-distance running? Eur J Sport Sci. 2016;16(8):965-71. Epub 2016/07/29. doi: 10.1080/17461391.2016.1200674. PubMed PMID: 27467184.
- 73. Brown N, Burnett E, Scurr J. Is Breast Pain Greater in Active Females Compared to the General Population in the UK? Breast J. 2016;22(2):194-201. Epub 2015/12/15. doi: 10.1111/tbj.12547. PubMed PMID: 26661830.
- Risius D, Milligan A, Berns J, Brown N, Scurr J.
 Understanding key performance indicators for
 breast support: An analysis of breast support effects
 on biomechanical, physiological and subjective
 measures during running. J Sports Sci.
 2017;35(9):842-51. Epub 2016/06/14. doi:
 10.1080/02640414.2016.1194523. PubMed PMID:
 27291899.
- 75. Burbage J, Cameron L. An investigation into the prevalence and impact of breast pain, bra issues and breast size on female horse riders. J Sports Sci. 2017;35(11):1091-7. Epub 2016/07/28. doi: 10.1080/02640414.2016.1210818. PubMed PMID: 27454814.
- Coltman CE, Steele JR, McGhee DE. Can breast characteristics predict upper torso musculoskeletal pain? Clin Biomech (Bristol, Avon). 2018;53:46-53. Epub 2018/02/16. doi: 10.1016/j.clinbiomech.2018.02.002. PubMed PMID: 29448080.
- 77. McGhee DE, Coltman KA, Riddiford-Harland DL, Steele JR. Upper torso pain and musculoskeletal structure and function in women with and without large breasts: A cross sectional study. Clin Biomech (Bristol, Avon). 2018;51:99-104. Epub 2017/12/30. doi: 10.1016/j.clinbiomech.2017.12.009. PubMed PMID: 29287172.
- 78. Steele JR, Gho SA, Campbell TE, Richards CJ, Beirne S, Spinks GM, et al. The Bionic Bra: Using electromaterials to sense and modify breast support to enhance active living. J Rehabil Assist Technol Eng. 2018;5:2055668318775905. Epub 2019/06/14. doi: 10.1177/2055668318775905. PubMed PMID: 31191941; PubMed Central PMCID: PMCPMC6453067.
- 79. Brown N, Smith J, Brasher A, Risius D, Marczyk A, Wakefield-Scurr J. Breast education for schoolgirls; why, what, when, and how? Breast J. 2018;24(3):377-82. Epub 2017/11/16. doi: 10.1111/tbj.12945. PubMed PMID: 29139615.
- 80. Smith LJ, Eichelberger TD, Kane EJ. Breast Injuries in Female Collegiate Basketball, Soccer, Softball and Volleyball Athletes: Prevalence, Type and Impact on Sports Participation. Eur J Breast Health. 2018;14(1):46-50. Epub 2018/01/13. doi:



- 10.5152/ejbh.2017.3748. PubMed PMID: 29322119; PubMed Central PMCID: PMCPMC5758063.
- 81. Arch ES, Colon S, Richards JG. A Comprehensive Method to Measure 3-Dimensional Bra Motion During Physical Activity. J Appl Biomech. 2018;34(5):392-5. Epub 2018/05/29. doi: 10.1123/jab.2017-0111. PubMed PMID: 29809078.
- 82. Brisbine BR, Steele JR, Phillips EJ, McGhee DE. The Occurrence, Causes and Perceived Performance Effects of Breast Injuries in Elite Female Athletes. J Sports Sci Med. 2019;18(3):569-76. Epub 2019/08/21. PubMed PMID: 31427880; PubMed Central PMCID: PMCPMC6683617.
- Coltman CE, Steele JR, McGhee DE. Effect of Breast Size on Upper Torso Musculoskeletal Structure and Function: A Cross-Sectional Study. Plast Reconstr Surg. 2019;143(3):686-95. Epub 2019/03/01. doi: 10.1097/PRS.0000000000005319. PubMed PMID: 30817639.
- 84. Leme JC, Banks LDS, Reis YBD, Cardoso JR, Campos MH, Santiago PRP, et al. Sports bra but not sports footwear decreases breast movement during walking and running. J Biomech. 2020;111:110014. Epub 2020/09/07. doi: 10.1016/j.jbiomech.2020.110014. PubMed PMID: 32891811.
- 85. Leme JC, Reis YBD, Banks LDS, Cardoso JR, Campos MH, Moura FA. Effects of sports bra and footwear on vertebral posture during walking and running. J Biomech. 2020;99:109524. Epub 2019/12/04. doi: 10.1016/j.jbiomech.2019.109524. PubMed PMID: 31787256.
- 86. Norris M, Mills C, Sanchez A, Wakefield-Scurr J. Do static and dynamic activities induce potentially damaging breast skin strain? BMJ Open Sport Exerc Med. 2020;6(1):e000770. Epub 2020/07/24. doi: 10.1136/bmjsem-2020-000770. PubMed PMID: 32699646; PubMed Central PMCID: PMCPMC7365429.
- 87. Navalta JW, Ramirez GG, Maxwell C, Radzak KN, McGinnis GR. Validity and Reliability of Three Commercially Available Smart Sports Bras during Treadmill Walking and Running. Sci Rep. 2020;10(1):7397. Epub 2020/05/02. doi: 10.1038/s41598-020-64185-z. PubMed PMID: 32355249; PubMed Central PMCID: PMCPMC7192924.
- 88. Brisbine BR, Steele JR, Phillips EJ, McGhee DE. Can Physical Characteristics and Sports Bra Use Predict Exercise-Induced Breast Pain in Elite Female Athletes? Clin J Sport Med. 2021;31(6):e380-e4. Epub 2020/02/20. doi: 10.1097/JSM.00000000000000831. PubMed PMID: 32073475.
- 89. Norris M, Blackmore T, Horler B, Wakefield-Scurr J. How the characteristics of sports bras affect their performance. Ergonomics. 2021;64(3):410-25. Epub 2020/09/29. doi: 10.1080/00140139.2020.1829090. PubMed PMID: 32981459.

- Jones M, Mills C, Exell T, Wakefield-Scurr J. A novel multi-study intervention investigating the short and long term effects of a posture bra on whole body and breast kinematics. Gait Posture. 2021;83:194-200. Epub 2020/11/09. doi: 10.1016/j.gaitpost.2020.10.031. PubMed PMID: 33161276.
- 91. Exell TA, Milligan A, Burbage J, Risius D, Sanchez A, Horler B, et al. There are two sides to every story: implications of asymmetry on breast support requirements for sports bra manufacturers. Sports Biomech. 2021;20(7):866-78. Epub 2019/06/15. doi: 10.1080/14763141.2019.1614654. PubMed PMID: 31198100.
- 92. Can TS, Ikizceli TU. A Rare Complication of The Heart Rate Strap Use in A Marathon Runner: Mondor's Disease of the Breast. J Coll Physicians Surg Pak. 2022;32(12):1623-5. Epub 2022/12/08. doi: 10.29271/jcpsp.2022.12.1623. PubMed PMID: 36474389.
- Haworth L, May K, Janssen J, Selfe J, Chohan A. The impact of breast support garments on fit, support and posture of larger breasted women. Appl Ergon. 2022;101:103701. Epub 2022/02/13. doi: 10.1016/j.apergo.2022.103701. PubMed PMID: 35151120.
- Wakefield-Scurr J, Sanchez A, Jones M, Hockley L, Biswas A, Johnson F, et al. A multi-phase intervention study of sports bra prescription for elite UK female athletes preparing for the Tokyo Olympics and Paralympics. Res Sports Med. 2022:1-15. Epub 2022/06/18. doi: 10.1080/15438627.2022.2090254. PubMed PMID: 35711113
- Wakefield-Scurr J, Hamilton C, Reeves K, Jones M, Jones B. The effect of washing and wearing on sports bra function. Sports Biomech. 2022:1-11. Epub 2022/03/02. doi: 10.1080/14763141.2022.2046147. PubMed PMID: 35227159.
- 96. Fong HB, Powell DW. Greater Breast Support Is Associated With Reduced Oxygen Consumption and Greater Running Economy During a Treadmill Running Task. Front Sports Act Living. 2022;4:902276. Epub 2022/07/02. doi: 10.3389/fspor.2022.902276. PubMed PMID: 35774380; PubMed Central PMCID: PMCPMC9237383.
- 97. Fong HB, Nelson AK, Storey JE, Hinton J, Puppa M, McGhee D, et al. Greater Breast Support Alters Trunk and Knee Joint Biomechanics Commonly Associated With Anterior Cruciate Ligament Injury. Front Sports Act Living. 2022;4:861553. Epub 2022/06/08. doi: 10.3389/fspor.2022.861553. PubMed PMID: 35669558; PubMed Central PMCID: PMCPMC9163541.
- 98. Rotllan C, Viscor G. Winter Sports Injuries in Elite Female Athletes: A Narrative Review. Int J Environ Res Public Health. 2023;20(10). Epub 2023/05/27.



- doi: 10.3390/ijerph20105815. PubMed PMID: 37239544; PubMed Central PMCID: PMCPMC10218179.
- 99. Mullins K, Mac Colgain D, Carton P. Incidence and Severity of Hamstring Injuries in Female Athletes Who Play Field Sports: A Systematic Review With Meta-Analysis of Prospective Studies. J Orthop Sports Phys Ther. 2022;52(11):740-A5. Epub 2022/08/06. doi: 10.2519/jospt.2022.11144. PubMed PMID: 35930061.
- 100. Petushek EJ, Sugimoto D, Stoolmiller M, Smith G, Myer GD. Evidence-Based Best-Practice Guidelines for Preventing Anterior Cruciate Ligament Injuries in Young Female Athletes: A Systematic Review and Meta-analysis. Am J Sports Med. 2019;47(7):1744-53. Epub 2018/07/13. doi: 10.1177/0363546518782460. PubMed PMID: 30001501; PubMed Central PMCID: PMCPMC6592422.
- 101. Crossley KM, Patterson BE, Culvenor AG, Bruder AM, Mosler AB, Mentiplay BF. Making football safer for women: a systematic review and meta-analysis of injury prevention programmes in 11 773 female football (soccer) players. Br J Sports Med. 2020;54(18):1089-98. Epub 2020/04/08. doi: 10.1136/bjsports-2019-101587. PubMed PMID: 32253193; PubMed Central PMCID: PMCPMC7497572.
- 102. Lopez-Valenciano A, Raya-Gonzalez J, Garcia-Gomez JA, Aparicio-Sarmiento A, Sainz de Baranda P, De Ste Croix M, et al. Injury Profile in Women's Football: A Systematic Review and Meta-Analysis. Sports Med. 2021;51(3):423-42. Epub 2021/01/13. doi: 10.1007/s40279-020-01401-w. PubMed PMID: 33433863.
- 103. Maya J, Misra M. The female athlete triad: review of current literature. Curr Opin Endocrinol Diabetes Obes. 2022;29(1):44-51. Epub 2021/11/24. doi: 10.1097/MED.0000000000000690. PubMed PMID: 34812202; PubMed Central PMCID: PMCPMC8702454.
- 104. Thein-Nissenbaum J, Hammer E. Treatment strategies for the female athlete triad in the adolescent athlete: current perspectives. Open Access J Sports Med. 2017;8:85-95. Epub 2017/04/25. doi: 10.2147/OAJSM.S100026. PubMed PMID: 28435337; PubMed Central PMCID: PMCPMC5388220.
- 105. de Borja C, Chang CJ, Watkins R, Senter C. Optimizing Health and Athletic Performance for Women. Curr Rev Musculoskelet Med. 2022;15(1):10-20. Epub 2022/01/14. doi: 10.1007/s12178-021-09735-2. PubMed PMID: 35023069; PubMed Central PMCID: PMCPMC8804053.
- 106. Randell RK, Clifford T, Drust B, Moss SL, Unnithan VB, De Ste Croix MBA, et al. Physiological Characteristics of Female Soccer Players and Health and Performance Considerations: A Narrative

- Review. Sports Med. 2021;51(7):1377-99. Epub 2021/04/13. doi: 10.1007/s40279-021-01458-1. PubMed PMID: 33844195; PubMed Central PMCID: PMCPMC8222040 and Daniel Martin declare that they have no conflicts of interest relevant to the content of this review. Rebecca K Randell, James M Carter and Ian Rollo are employees of the Gatorade Sports Science Institute, a division of PepsiCo, Incorporated. Samantha L Moss is a paid consultant for the Gatorade Sports Science Institute. The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of PepsiCo, Incorporated. Hannah Mayho is an employee of Manchester City Football Club.
- 107. Elliott-Sale KJ, McNulty KL, Ansdell P, Goodall S, Hicks KM, Thomas K, et al. The Effects of Oral Contraceptives on Exercise Performance in Women: A Systematic Review and Meta-analysis. Sports Med. 2020;50(10):1785-812. Epub 2020/07/16. doi: 10.1007/s40279-020-01317-5. PubMed PMID: 32666247; PubMed Central PMCID: PMCPMC7497464 Kevin Thomas, Paul Swinton and Eimear Dolan declare that they have no potential conflicts of interest with the content of this article.
- 108. Carmichael MA, Thomson RL, Moran LJ, Wycherley TP. The Impact of Menstrual Cycle Phase on Athletes' Performance: A Narrative Review. Int J Environ Res Public Health. 2021;18(4). Epub 2021/02/13. doi: 10.3390/ijerph18041667. PubMed PMID: 33572406; PubMed Central PMCID: PMCPMC7916245.
- 109. Holtzman B, Ackerman KE. Recommendations and Nutritional Considerations for Female Athletes: Health and Performance. Sports Med. 2021;51(Suppl 1):43-57. Epub 2021/09/14. doi: 10.1007/s40279-021-01508-8. PubMed PMID: 34515972; PubMed Central PMCID: PMCPMC8566643 the content of this article.
- 110. McGhee DE, Steele JR. Breast Biomechanics: What Do We Really Know? Physiology (Bethesda). 2020;35(2):144-56. Epub 2020/02/07. doi: 10.1152/physiol.00024.2019. PubMed PMID: 32027563.
- 111. Zhou J, Yu W, Ng S-P. Methods of studying breast motion in sports bras: a review. Textile Research Journal. 2011;81(12):1234-48. doi: 10.1177/0040517511399959.
- 112. Haworth L, Aitkenhead R, Grecic D, Chohan A. Understanding experience, knowledge and perceived challenges related to bra fit for sports participation: a scoping review. Res Sports Med. 2022:1-14. Epub 2022/07/21. doi: 10.1080/15438627.2022.2102922. PubMed PMID: 35856556.
- 113. Paul RW, Sonnier JH, Johnson EE, Hall AT, Osman A, Connors GM, et al. Inequalities in the Evaluation of Male Versus Female Athletes in Sports Medicine Research: A Systematic Review. Am J Sports Med. 2022;3635465221131281. Epub 2022/12/02. doi:



- 10.1177/03635465221131281. PubMed PMID: 36453705.
- 114. Murrar S, Johnson PA, Lee YG, Carnes M. Research Conducted in Women Was Deemed More Impactful but Less Publishable than the Same Research Conducted in Men. J Womens Health (Larchmt). 2021;30(9):1259-67. Epub 2021/03/16. doi: 10.1089/jwh.2020.8666. PubMed PMID: 33719578; PubMed Central PMCID: PMCPMC8561742.
- 115. Brisbine BR, Steele JR, Phillips EJ, McGhee DE. Breast pain affects the performance of elite female athletes. J Sports Sci. 2020;38(5):528-33. Epub 2020/01/11. doi: 10.1080/02640414.2020.1712016. PubMed PMID: 31918624.
- 116. Hadi MS. Sports Brassiere: Is It a Solution for Mastalgia? Breast J. 2000;6(6):407-9. Epub 2001/05/12. doi: 10.1046/j.1524-4741.2000.20018.x. PubMed PMID: 11348400.

- 117. Sport Wi. Only a third of secondary school girls wear a sports bra for PE 2023 [cited 2023 November 23]. Available from: https://womeninsport.org/news/only-a-third-of-
- 118. O'Sullivan S. Lack Of Sports Bras Sidelines Seattle Girls, But It Doesn't Have To. The Seattle Medium.

secondary-school-girls-wear-a-sports-bra-for-pe/.

- 119. LACKE S. Why Girls Stop Running in the Tween Years. triathlete. 2022.
- 120. Lewis S. 'How did none of us think of this before?': sports bra scheme breaks down entry barriers. The Guardian. 2020.
- 121. The Sports Bra Project 2023. Available from: https://www.thesportsbraproject.org/.
- 122. Bras for Girls 2023. Available from: https://www.brasforgirls.org/.